arXiv:quant-ph/0604003v1 1 Apr 2006

# ATOM-PHOTON INTERACTIONS WITH RESPECT TO QUANTUM COMPUTATION: THREE-LEVEL ATOM IN TWO-MODE FIELD

G. K. GIORGADZE AND Z. G. MELIKISHVILI

Abstract. In the paper, analysis of a quantum optical system—three-level atom in a quantum electromagnetic field is given. Evolution operators are constructed in closed form.

## 1. Introduction

In theory of computation it is known that the three-state element is most efficient for classical computers, but its realization remained a problem for many years. The progress was achieved in 90-ties (see [6], [7], [8]). Optimal number of states of the physical system for the expression mod $p$, where $p$ is a prime, is 3. Indeed, if we denote by $N_p$ this number, it is known that $N_p = f(p)N_2$, where $f(p) = \frac{p}{2\log_2 p}$, we obtain that the function $\frac{N_p}{N_2}$ have the minimum when $p = 3$. The necessary mathematical theory in classical many-valued logic is given in [9].

In the quantum world the many-level systems are ordinary ones. Thus in this paper we will consider one possible version of a quantum processor based on a many-level quantum system.

In [10] the extension of the universal quantum logic to the multi-valued domain is considered, where the unit of memory is the qudit. Thereby it is assumed that arbitrary unitary operators on any $d$-level system, where $d > 2$ is any number, can be decomposed into logical gates that operate on only two levels at a time. For example, the linear ion trap quantum computer [11] uses only two levels in each ion for computing although additional levels can be accessed, and are typically needed, for processing and reading out the state of the ion (see [10] and the literature cited there).

In the case when $d = 3$, the set of universal gates is built in the following manner.

Let us introduce the following unitary operators on $C^3 \to C^3$ :

$$X_3(\phi) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & e^{i\phi} \end{pmatrix}, Z_3(c_0, c_1, c_2) = \begin{pmatrix} c_0 & 0 & -c_2 \\ 0 & -c_1 & c_2 \\ -\overline{c}_2 & \overline{c}_2 & \overline{c}_2 \end{pmatrix}.$$

It can be easily checked that $X_3(\phi)$ and $Z_3(c_0, c_1, c_2)$ act on the states $|0>$, $|1>$, $|2>$ as

$$X_3(\phi)|0>= |0>, X_3(\phi)|1>= |1>, X_3(\phi)|2>= e^{i\phi}|2>;$$

$$Z_3(\phi)|0>= c_0|0> -\overline{c}_2|2>, Z_3(\phi)|1>= -c_1|0> +\overline{c}_2|2>, Z_3|2>= -c_2|0> +c_2|1> +\overline{c}_2|2> .$$

Furthermore, the gate $Z_3(c_0, c_1, c_2)$ satisfies the following condition:

$$Z_3(c_0, c_1, c_2) = \begin{pmatrix} c_0 & 0 & -c_2 \\ 0 & -c_1 & c_2 \\ -\overline{c}_2 & \overline{c}_2 & \overline{c}_2 \end{pmatrix} (c_0|0> +c_1|1> +c_2|2>) =$$

$$= (-c_2c_0 + c_0c_2, -c_2c_1 + c_1c_2, |c_0|^2 + |c_1|^2 + |c_2|^2 = |2>).$$

In [10] it is proved that the set of gates $\{X_3(\phi), Z_3(c_0, c_1, c_2), G_2[X_3(\phi)], G_2[Z_3(c_0, c)1, c_2)]\}$ is universal for a three-level quantum system, where $G_2[Y] = \begin{pmatrix} I & 0 \\ 0 & Y \end{pmatrix}$ and $I$ is the identity matrix.

Below is given another approach of description of quantum system which gives the single qutrit gates, in particular, we obtain in closed form evolution operators depending on parameters.

---

To appear in “Contemporary mathematics and its applications”.

## 2. ATOM-PHOTON INTERACTIONS HAMILTONIAN

We will build the special case Hamiltonian of a hydrogen-like atom moving and interacting with a quantized electromagnetic field, at that using the notations of [1] and [3]. The general form of such Hamiltonian, including the center-of-mass motion and all the modes of the interacting field, is presented in [1]. For our purposes we consider the following conditions on atom-photon interactions: (i) All the atom-photon interactions are the electric dipole; (ii) Only three atomic levels are included in the interaction; (iii) Two quantized resonator modes are interacting with this three-level system; (iv) Each of these modes interacts with only one couple of levels; (v) So, from three possible couples of levels, only two of them interact directly. Levels of the third couple interact only by means of an intermediate level; (vi) In this work we permit, but do not consider the center-of-mass motion and large detuning from resonances, which cause the state entanglement.

Let us consider atom-photon interactions Hamiltonian in the electric dipole approximation

$$\hat{H}_{\mathbf{r}.\mathbf{E}} = -\wp\mathbf{E}(R,t) = -e\mathbf{r}\mathbf{E}(\mathbf{R},t), \tag{1}$$

and transform it for three-level atom interacting with two modes of quantized fields of radiation. For these purposes previously we must discuss the following operators.

**Hamiltonian for internal degrees of freedom of non-interacting atom.** We consider internal degrees of freedom of a three-level atom. Due to assumptions (i-vi) only two couples of energy states are involved in interaction. In all cases the intermediate state with energy $E_i \equiv \hbar\omega_i$ is labeled by the state vector $|i>$. From other states, the state with higher energy is named excited, denoted $E_e \equiv \hbar\omega_e$ and labeled by the state vector $|e>$. The remaining energy level is named ground level, denoted $E_g \equiv \hbar\omega_g$ and labeled by the state vector $|g>$ (see Fig.1). All of them are eigenstates of the Hamiltonian $\hat{H}_{atom}$ of non-interacting three-level atom:

$$\hat{H}_{atom}|e>=\hbar\omega_e|e>,\ \hat{H}_{atom}|i>=\hbar\omega_i|i>,\ \hat{H}_{atom}|g>=\hbar\omega_g|g>. \tag{2}$$

Using the completeness relation $|e><e|+|i><i|+|g><g|=1$, we can write the arbitrary atomic operator $\hat{\mathbf{O}}$ as

$$\hat{\mathbf{O}} = 1\hat{\mathbf{O}}1 = \sum_{j,j'=e,g,i} |j><j|\hat{\mathbf{O}}|j'><j'|. \tag{3}$$

Utilizing (3) and the orthonormality condition $<j||j'>=\delta_{ij'}$ for the Hamiltonian of a three-level atom, one can easily obtain

$$\begin{aligned}\hat{H}_{atom} &= (|e><e|+|i><i|+|g><g|)\hat{H}_{atom}(|e><e|+|i><i|+|g><g|) = \\ &E_e|e><e|+E_i|i><i|+E_g|g><g|,\end{aligned} \tag{4}$$

where we use (2) and above notations $E_e=\hbar\omega_e$, $E_i=\hbar\omega_i$, $E_g=\hbar\omega_g$. Then

$$|e>=\begin{pmatrix}1\\0\\0\end{pmatrix}, <e|=(1,0,0), |i>=\begin{pmatrix}0\\1\\0\end{pmatrix}, <i|=(0,1,0), |g>=\begin{pmatrix}0\\0\\1\end{pmatrix}, <g|=(0,0,1),$$

Thus, the Hamiltonian (4) has the following matrix form

$$\begin{aligned}\hat{H}_{atom} &= E_e\begin{pmatrix}1&0&0\\0&0&0\\0&0&0\end{pmatrix} + E_i\begin{pmatrix}0&0&0\\0&1&0\\0&0&0\end{pmatrix} + E_g\begin{pmatrix}0&0&0\\0&0&0\\0&0&1\end{pmatrix} = \\ &= \begin{pmatrix}E_g&0&0\\0&E_i&0\\0&0&E_g\end{pmatrix} = \hbar\begin{pmatrix}\omega_e&0&0\\0&\omega_i&0\\0&0&\omega_g\end{pmatrix}.\end{aligned} \tag{5}$$

**The dipole moment operator.** Now we can use the completeness relation once more to express the position operator $\hat{\mathbf{r}}$ in terms of energy eigenstates. As each state, wave functions $\Psi_j(\mathbf{r})$ of a free atom have definite parity, then the diagonal matrix elements equal zero

$$<j|\hat{\mathbf{r}}|j>=\int|\Psi_j(\mathbf{r})|^2\mathbf{r}d^3r=0.$$

Indeed, if the function $|\Psi_j(\mathbf{r})|^2$ is symmetric and the operator $\mathbf{r}$ antisymmetric, the expression under the integral sign is antisymmetric. For the nondiagonal matrix elements we have the following expressions

$$e < i|\hat{\mathbf{r}}|g >= e \int \Psi_i^*(\mathbf{r})\mathbf{r}\Psi_g(\mathbf{r})d^3r = \wp_{ig}, \tag{6}$$

and

$$e < g|\hat{\mathbf{r}}|i >= e \int \Psi_g^*(\mathbf{r})\mathbf{r}\Psi_i(\mathbf{r})d^3r = \wp_{ig}^* = \wp_{gi}.$$

In the same way

$$e < e|\hat{\mathbf{r}}|i >= e \int \Psi_i^*(\mathbf{r})\mathbf{r}\Psi_g(\mathbf{r})d^3r = \wp_{ei},$$

and

$$e < i|\hat{\mathbf{r}}|e >= e \int \Psi_g^*(\mathbf{r})\mathbf{r}\Psi_i(\mathbf{r})d^3r = \wp_{ei}^* = \wp_{ie}.$$

Thus, the dipole moment operator $e\hat{\mathbf{r}}$ is expressed in the following manner

$$e\hat{\mathbf{r}} = \wp_{ig}|i >< g| + \wp_{gi}|g >< i| + \wp_{ei}|e >< i| + \wp_{ie}|i >< e|. \tag{7}$$

Note that this operator describes transitions from the ground state $|g >$ to the intermediate state $|i >$, from intermediate to the excited state $|e >$, and the process in reverse order back to $|g >$. For verification let uss apply the operator $e\hat{\mathbf{r}}$ (9) to the ground state $|g >$ . Then we get

$$e\hat{\mathbf{r}}|g >= \wp_{ig}|i >< g||g > +\wp_{gi}|g >< i||g > +\wp_{ei}|e >< i||g > +\wp_{ie}|i >< e||g >= \wp_{ig}|i > .$$

Similarly

$$e\hat{\mathbf{r}}|e >= \wp_{ig}|i >< g||e > +\wp_{gi}|g >< i||e > +\wp_{ei}|e >< i||e > +\wp_{ie}|i >< e||e >= \wp_{ie}|i >$$

and finally

$$e\hat{\mathbf{r}}|i >= \wp_{ig}|i >< g||i > +\wp_{gi}|g >< i||i > +\wp_{ei}|e >< i||i > +\wp_{ie}|i >< e||g >= \wp_{gi}|g > +\wp_{ei}|e > .$$

Thus, the operator

$$\hat{\sigma}_{ig} \equiv |i >< g|,\ |i >< g| = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} (0,0,1) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}, \tag{8}$$

causes the atom transition from the ground to the intermediate state and can be interpreted as a creation operator for the atom in the intermediate state $|i >$.

On the contrary, the operator

$$\hat{\sigma}_{gi} \equiv |g >< i|,\ |g >< i| = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} (0,1,0) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \tag{9}$$

annihilates the atom in the intermediate state and so represents the annihilation operator for the intermediate state.

The operator

$$\hat{\sigma}_{ei} \equiv |e >< i|,\ |e >< i| = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} (0,1,0) = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \tag{10}$$

causes transition of the atom from the intermediate to the excited state and can be interpreted as a creation operator for the atom in the excited state $|e >$.

The operator

$$\hat{\sigma}_{ei} \equiv |i >< e|,\ |i >< e| = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} (1,0,0) = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \tag{11}$$

annihilates the atom in the excited state and so represents the annihilation operator for the excited state. The dipole moment operator (7) is defined in accordance to the expressions (8)-(11), of creation and annihilation operators, to be

$$e\hat{\mathbf{r}} = \wp_{ig}\hat{\sigma}_{ig} + \wp_{gi}\hat{\sigma}_{ig} + \wp_{ei}\hat{\sigma}_{ei} + \wp_{ie}\hat{\sigma}_{ie}. \tag{12}$$

**The electric field operator.** Now let us discuss two modes of radiation field. We assume for determinacy, that the radiation field is formed in three-mirror standing wave resonator and decompose the electric field strength in the resonator as follows

$$\hat{\mathbf{E}}(\hat{\mathbf{R}},t) = \mathcal{F}_{0a}\mathbf{u}_a(\hat{\mathbf{R}})i(\hat{a}-\hat{a}^+) + \mathcal{F}_{0b}\mathbf{u}_b(\hat{\mathbf{R}})i(\hat{b}-\hat{b}^+), \tag{13}$$

where $\mathcal{F}_{0j} \equiv \sqrt{\frac{\hbar\Omega_j}{2\epsilon_0 V_j}}$, $j = a, b$.

Here $\mathbf{u}_j(\hat{\mathbf{R}})$ represents the mode functions of the resonator at the position $\mathbf{R}$ of the atom. $\Omega_a$ and $\Omega_b$ are oscillation frequencies of the field. In a classical treatment, $\hat{a}^+$, $\hat{b}^+$ and $\hat{a}$, $\hat{b}$ are time dependent amplitudes. In our present treatment they are operators which obey the commutation relations

$$\hat{a}\hat{a}^+ - \hat{a}^+\hat{a} = 1,\ \hat{b}\hat{b}^+ - \hat{b}^+\hat{b} = 1. \tag{14}$$

With their aid the quantized field Hamiltonian can be written in the form

$$\hat{H}_{field} = \hbar\Omega_a\hat{a}^+\hat{a} + \hbar\Omega_b\hat{b}^+\hat{b}. \tag{15}$$

Here we neglect the zero-point energy.

**Back to the interaction Hamiltonian.** Now consider the Hamiltonian of electric dipole atom-photon interactions again (1)

$$\hat{H}_{\mathbf{rE}} = -e\hat{\mathbf{r}}\hat{\mathbf{E}}(\hat{\mathbf{R}},t).$$

Substituting in this expression the expressions of the dipole moment operator $e\hat{\mathbf{r}}$ (12) and electric field operator $\hat{\mathbf{r}}\hat{\mathbf{E}}(\hat{\mathbf{R}},t)$ we obtain

$$\begin{aligned}\hat{H}_{\mathbf{rE}} = &-\mathcal{F}_{0a}i[\wp_{ig}\mathbf{u}_a(\hat{\mathbf{R}})\hat{\sigma}_{ig} + \wp_{ei}\mathbf{u}_b(\hat{\mathbf{R}})\hat{\sigma}_{gi}](\hat{a}-\hat{a}^+)- \\ &-\mathcal{F}_{0b}i[\wp_{ei}\mathbf{u}_b(\hat{\mathbf{R}})\hat{\sigma}_{ei} + \wp_{ei}\mathbf{u}_b(\hat{\mathbf{R}})\hat{\sigma}_{ie}](\hat{b}-\hat{b}^+).\end{aligned} \tag{16}$$

This expression contains the scalar product $\wp\mathbf{u}$ of the dipole moment and the mode function $\mathbf{u}$. As this product is a complex quantity, we represent it in the following manner

$$\wp\mathbf{u} = |\wp\mathbf{u}|e^{i\varphi}, \tag{17}$$

where $\varphi$ is the phase. Substituting (17) in (16) the Hamiltonian of electric dipole atom-photon interactions can be written in the following form

$$\begin{aligned}\hat{H}_{\mathbf{rE}} = &-\hbar\mathcal{F}_{0a}i[\frac{|\wp_{ig}\mathbf{u}_a(\hat{\mathbf{R}})|}{\hbar}\hat{\sigma}_{ig}e^{i\varphi_a} + \frac{|\wp_{ei}\mathbf{u}_b(\hat{\mathbf{R}})|}{\hbar}\hat{\sigma}_{gi}e^{-i\varphi_a}](\hat{a}-\hat{a}^+)- \\ &-\hbar\mathcal{F}_{0b}i[\frac{|\wp_{ei}\mathbf{u}_b(\hat{\mathbf{R}})|}{\hbar}\hat{\sigma}_{ei}e^{i\varphi_b} + \frac{|\wp_{ei}\mathbf{u}_b(\hat{\mathbf{R}})|}{\hbar}\hat{\sigma}_{ie}e^{-i\varphi_b}](\hat{b}-\hat{b}^+).\end{aligned} \tag{18}$$

Introducing the Rabi frequencies

$$g_a(\mathbf{R}) = \frac{|\wp_{ig}\mathbf{u}_a(\hat{\mathbf{R}})|}{\hbar}\mathcal{F}_{0a},\ g_b(\mathbf{R}) = \frac{|\wp_{ei}\mathbf{u}_b(\hat{\mathbf{R}})|}{\hbar}\mathcal{F}_{0b}$$

the interaction Hamiltonian (18) now is given by

$$\begin{aligned}\hat{H}_{\mathbf{rE}} = &\hbar g_a(\hat{\mathbf{R}})(-i)(\hat{\sigma}_{ig}e^{i\varphi_a} + \hat{\sigma}_{gi}e^{-i\varphi_a})(\hat{a}-\hat{a}^+)+ \\ &\hbar g_b(\hat{\mathbf{R}})(-i)(\hat{\sigma}_{ei}e^{i\varphi_b} + \hat{\sigma}_{ie}e^{-i\varphi_b})(\hat{b}-\hat{b}^+).\end{aligned} \tag{19}$$

Introducing the difference of phases $\delta\varphi \equiv \varphi_a - \varphi_b$, in the Hamiltonian (19) we obtain

$$\begin{aligned}\hat{H}_{\mathbf{rE}} = &\hbar g_a(\hat{\mathbf{R}})(-i)(\hat{\sigma}_{ig}e^{i\varphi_a} + \hat{\sigma}_{gi}e^{-i\varphi_a})(\hat{a}-\hat{a}^+)+ \\ &\hbar g_b(\hat{\mathbf{R}})(-i)(\hat{\sigma}_{ei}e^{i(\delta\varphi-\varphi_a)} + \hat{\sigma}_{ie}e^{-i(\delta\varphi-\varphi_a)})(\hat{b}-\hat{b}^+).\end{aligned} \tag{20}$$

In quantum mechanics only the difference of phases is important, so in (20) we can put $\varphi_a = \pi/2$ and finally obtain

$$\hat{H}_{\mathbf{rE}} = \hbar g_a(\hat{\mathbf{R}})(\hat{\sigma}_{ig} - \hat{\sigma}_{gi})(\hat{a}-\hat{a}^+) + \hbar g_b(\hat{\mathbf{R}})(\hat{\sigma}_{ei}e^{-i\delta\varphi} - \hat{\sigma}_{ie}e^{i\delta\varphi})(\hat{b}-\hat{b}^+). \tag{21}$$

Expression (21) is the final form of the exact Hamiltonian responsible for all atom-photon interactions between the three-level atom and two modes of electromagnetic field. Just this Hamiltonian will be used hereinafter.

**Total Hamiltonian.** Joining up the above main results (4), (15), (21), and taking into account, that

$$\hat{\sigma}_{ee} \equiv |e><e|,\ \hat{\sigma}_{ii} \equiv |i><i|,\ \hat{\sigma}_{gg} \equiv |g><g|,$$

one can build the total Hamiltonian of the whole atom-field system

$$\hat{H} = \frac{\hat{\mathbf{P}}^2}{2M} + \hat{H}_{field} + \hat{H}_{atom} + \hat{H}_{\mathbf{rE}} = \tag{22}$$

$$\frac{\hat{\mathbf{P}}^2}{2M} + \hbar\Omega_a \hat{a}^+\hat{a} + \hbar\Omega_b \hat{b}^+\hat{b} + E_e\hat{\sigma}_{ee} + E_i\hat{\sigma}_{ii} + E_g\hat{\sigma}_{gg} +$$

$$+\hbar g_a(\hat{\mathbf{R}})(\hat{\sigma}_{ig} - \hat{\sigma}_{gi})(\hat{a} - \hat{a}^+) + \hbar g_b(\hat{\mathbf{R}})(\hat{\sigma}_{ei}e^{-i\delta\varphi} - \hat{\sigma}_{ie}e^{i\delta\varphi})(\hat{b} - \hat{b}^+).$$

This Hamiltonian includes: the operator of kinetic energy $\frac{\hat{\mathbf{P}}^2}{2M}$ of the center of mass movement; free-field Hamiltonian $\hat{H}_{field} = \hbar\Omega_a\hat{a}^+\hat{a} + \hbar\Omega_b\hat{b}^+\hat{b}$; the Hamiltonian responsible for internal states of the atom $\hat{H}_{atom} = E_e\hat{\sigma}_{ee} + E_i\hat{\sigma}_{ii} + E_g\hat{\sigma}_{gg}$ and interaction operator $\hat{H}_{\mathbf{rE}} = \hbar g_a(\hat{\mathbf{R}})(\hat{\sigma}_{ig} - \hat{\sigma}_{gi})(\hat{a} - \hat{a}^+) + \hbar g_b(\hat{\mathbf{R}})(\hat{\sigma}_{ei}e^{-i\delta\varphi} - \hat{\sigma}_{ie}e^{i\delta\varphi})(\hat{b} - \hat{b}^+)$. Connecting the internal states with field operators $\hat{a} - \hat{a}^+$, $\hat{b} - \hat{b}^+$ by means of atom-field coupling constants $g_a(\hat{\mathbf{R}})$, $g_a(\hat{\mathbf{R}})$.

## 3. SIMPLIFICATION OF ATOM-PHOTON INTERACTIONS HAMILTONIAN BY SELECTING THE REAL TRANSITIONS (SRT) AND ROTATING WAVE APPROXIMATION (RWA)

There are two reasons for simplification of the atom-photon interactions Hamiltonian (21)

$$\hat{H}_{\mathbf{rE}} = \hbar g_a(\hat{\mathbf{R}})(\hat{\sigma}_{ig}\hat{a}^+ + \hat{\sigma}_{gi}\hat{a} - \hat{\sigma}_{ig}\hat{a} - \hat{\sigma}_{ig}\hat{a}^+) + \tag{23}$$

$$+\hbar g_b(\hat{\mathbf{R}})(\hat{\sigma}_{ei}\hat{b}^+e^{i\delta\varphi} - \hat{\sigma}_{ie}\hat{b}e^{-i\delta\varphi} - \hat{\sigma}_{ie}\hat{b}e^{i\delta\varphi} - \hat{\sigma}_{ie}\hat{b}^+e^{-i\delta\varphi}).$$

Namely, it amounts to confining our analysis to real transitions in which a photon is emitted while the atom goes from its upper state to its lower state, or a photon is absorbed while the atom goes from its lower state to its upper state. The second reason is mathematical and is connected with the averaging procedure – neglecting the fast oscillating terms. Hereinafter we will use the interaction representation (Dirac representation) which is defined by the internal atomic states and the free-field states and under these assumptions we may reduce the Hamiltonian (23) to the form which is called Rotating Wave Approximation (RWA).

In a three-level system the simplification procedure of interaction Hamiltonian should be carried out separately for each transition scheme. Thus, first of all let us consider these transition schemes. In the three-level atom due to (i)-(v) of section 2 there are three different configurations possible for transitions between the three levels depending upon the position of the intermediate level $|i>$. Those are shown in Fig. 1 along with the allowed transitions. In the configuration of the first diagrams in Fig. 1, the energy of $|i>$ is intermediate between the energies of the other two levels. It is called the ladder configuration. Notation 'L' stands for this configuration. The energy of $|i>$ in the $\Lambda$ or Raman configuration is higher than that of the other two levels (second diagram in Fig. 1) whereas the level $|i>$ in the V-system is below the other two levels (third diagram in Fig. 1).


a) L-configuration

b) Λ-configuration

c) V-configuration

Figure 1. Three configuration of the three level atom in a two-mode field each acting on only one transition.

3.1. **Simplification for L-configuration - SRT. Selection of real transitions.** The terms $\hat{\sigma}_{gi}\hat{a}$, $\hat{\sigma}_{ig}\hat{a}^{+}$, $\hat{\sigma}_{ie}\hat{b}$ and $\hat{\sigma}_{ei}\hat{b}^{+}$, in the interaction Hamiltonian (23) lead to the violation of the energy conservation law. Indeed, note that the operators $\hat{\sigma}_{gi}\hat{a}$ and $\hat{\sigma}_{ie}\hat{b}$, annihilate the excited atom (in $|i>$ and $|e>$) and simultaneously annihilate the field excitation (Fig. 2c). In the same manner operators $\hat{\sigma}_{ig}\hat{a}^{+}$ and $\hat{\sigma}_{ei}\hat{b}^{+}$ create the field excitation while the atom is annihilated in $|g>$ ($|i>$) and created in $|i>$ ($|e>$) states (Fig. 2d).



Figure 2.

On the contrary, the operators $\hat{\sigma}_{gi}\hat{a}^{+}$, $\hat{\sigma}_{ie}\hat{b}^{+}$ and $\hat{\sigma}_{ig}\hat{a}^{+}$, $\hat{\sigma}_{ei}\hat{b}$ either annihilate the excitation ($|i>$ or $|e>$) while creating the field excitation (Fig. 2b), or create the excited atom (in $|i>$ or $|e>$) while annihilating the field excitation (Fig. 2a).

Under the assumptions of real transitions, neglecting the operators $\hat{\sigma}_{gi}\hat{a}$, $\hat{\sigma}_{ie}\hat{b}$, $\hat{\sigma}_{ig}\hat{a}^{+}$, $\hat{\sigma}_{ie}\hat{b}^{+}$, we may reduce the Hamiltonian (23) to the form

$$\hat{H}^{L}_{\mathbf{rE}} \cong \hat{H}^{L}_{int} = \hbar g_a(\hat{\mathbf{R}})(\hat{\sigma}_{gi}\hat{a}^{+} + \hat{\sigma}_{ig}\hat{a}) + \hbar g_b(\hat{\mathbf{R}})(\hat{\sigma}_{ie}\hat{b}^{+}e^{i\delta\varphi} + \hat{\sigma}_{ei}\hat{b}e^{-i\delta\varphi}). \tag{24}$$

3.2. **Simplification for Λ-configuration. Selection of real transitions.** For this configuration the operators $\hat{\sigma}_{gi}\hat{a}$, $\hat{\sigma}_{ig}\hat{a}^{+}$, $\hat{\sigma}_{ie}\hat{b}^{+}$, $\hat{\sigma}_{ei}\hat{b}$ of the interaction Hamiltonian lead to the violation of the energy conservation law (see Fig.1 and Fig. 3c,d). Thus, neglecting these terms in the interaction Hamiltonian (23) and keeping $\hat\sigma_{gi}\hat a^+$, $\hat\sigma_{ig}\hat a$, $\hat\sigma_{ie}\hat b$ and $\hat\sigma_{ei}\hat b^+$, and (Fig. 3a,b), we may reduce the Hamiltonian (23) to the form



Figure 3.

$$\hat H^{\Lambda}_{\mathbf{rE}} \cong \hat H^{L}_{int} = \hbar g_a(\hat{\mathbf R})(\hat\sigma_{gi}\hat a^+ + \hat\sigma_{ig}\hat a) - \hbar g_b(\hat{\mathbf R})(\hat\sigma_{ie}\hat b^+ e^{i\delta\varphi} + \hat\sigma_{ei}\hat b e^{-i\delta\varphi}). \tag{25}$$

3.3. **Simplification for V-configuration. Selection of real transitions.** For this configuration the operators $\hat\sigma_{gi}\hat a^+$, $\hat\sigma_{ig}\hat a$, $\hat\sigma_{ie}\hat b$, and $\hat\sigma_{ei}\hat b^+$, of the interaction Hamiltonian lead to the violation of the energy conservation law (see Fig.1 and Fig. 4c,d). Thus, neglecting these terms in the interaction Hamiltonian (23) and keeping $\hat\sigma_{gi}\hat a$, $\hat\sigma_{ig}\hat a$, $\hat\sigma_{ie}\hat b$, and $\hat\sigma_{ei}\hat b$, and (Fig. 4a,b), we may reduce the Hamiltonian (23) to the form



Figure 4.

$$\hat H^{\Lambda}_{\mathbf{rE}} \cong \hat H^{V}_{int} = -\hbar g_a(\hat{\mathbf R})(\hat\sigma_{gi}\hat a^+ + \hat\sigma_{ig}\hat a) + \hbar g_b(\hat{\mathbf R})(\hat\sigma_{ie}\hat b^+ e^{i\delta\varphi} + \hat\sigma_{ei}\hat b e^{-i\delta\varphi}). \tag{26}$$

3.4. **Interaction representation.** Further simplification of the atom-photon interaction hamiltonians (21)-(26) takes place in the interaction representation, where we can use the averaging procedure by fast oscillations. To accomplish this representation we should represent the total Hamiltonian of the whole atom-field system (see (22))

$$\hat H = \frac{\hat{\mathbf P}^2}{2M} + \hat H_{field} + \hat H_{atom} + \hat H_{\mathbf{rE}} =$$

$$\frac{\hat{\mathbf P}^2}{2M} + \hat H_0 + \hbar g_a(\hat{\mathbf R})(\hat\sigma_{ig} - \hat\sigma_{gi})(\hat a - \hat a^+) + \hbar g_b(\hat{\mathbf R})(\hat\sigma_{ei}e^{i\delta\varphi} - \hat\sigma_{ie}e^{-i\delta\varphi}(\hat b - \hat b^+).$$

by the internal states of the atom and free-field states. Here

$$\hat H_0 = \hat H_{fild} + \hat H_{atom} = \hbar\Omega_a\hat a^+\hat a + \hbar\Omega_b\hat b^+\hat b + E_e\hat\sigma_{ee} + E_i\hat\sigma_{ii} + E_g\hat\sigma_{gg},$$

or

$$\hat H_0 = \hbar\Omega_a\hat a^+\hat a + \hbar\Omega_b\hat b^+\hat b + \hbar\begin{pmatrix}\omega_e & 0 & 0\\ 0 & \omega_i & 0\\ 0 & 0 & \omega_g\end{pmatrix} \tag{27}$$

denotes the sum of free-field $\hat H_{field}$ and free atom $\hat H_{atom}$ Hamiltonians.

Now, defining the state vector in the interaction representation $|\Phi^{(I)}(t)>$ via

$$|\Phi(t)> \equiv exp[-\frac{i}{\hbar}\hat H_0 t]|\Phi^{(I)}(t)> \tag{28}$$

and substituting (28), containing the state vector $|\Phi^{(I)}(t)>$ in the interaction representation, in the Schrödinger equation

$$i\hbar\frac{d|\Phi^{(I)}(t)>}{dt} = (\frac{\hat{\mathbf{P}}^2}{2M} + \hat H_0 + \hat H_{\mathbf{rE}})|\Phi(t)> .$$

We receive

$$i\hbar\frac{d|\Phi^{(I)}(t)>}{dt} = (\frac{\hat{\mathbf{P}}^2}{2M} + \hat H^I_{\mathbf{rE}})|\Phi(t)>,$$

where

$$\hat H^{(I)}_{\mathbf{rE}} \equiv \exp[\frac{i}{\hbar}\hat H_0 t]\hat H_{\mathbf{rE}}\exp[-\frac{i}{\hbar}\hat H_0 t] \tag{29}$$

is the interaction Hamiltonian in the interaction picture. Note, that by reason of (vi) of section 2, we do not include the kinetic energy operator (corresponding to the center of mass movement) in above transformations. Now, let us obtain the exact expression for the interaction Hamiltonian $\hat H^{(I)}_{\mathbf{rE}}$. Substituting $\hat H_0$ (27) into the expression (29), under conditions, that the field and atom operators commute with each other we obtain

$$\begin{aligned} \hat H^{(I)}_{\mathbf{rE}} &= \hbar g_a(\hat{\mathbf{R}}) e^{\frac{i}{\hbar}\hat H_{atom}t}(\hat\sigma_{ig} - \hat\sigma_{gi})e^{-\frac{i}{\hbar}\hat H_{atom}t} e^{i\Omega_a\hat a^+\hat a t}(\hat a - \hat a^+)e^{-i\Omega_a\hat a^+\hat a t} \\ &+\hbar g_a(\hat{\mathbf{R}}) e^{\frac{i}{\hbar}\hat H_{atom}t}(\hat\sigma_{ei}e^{-i\delta\varphi} - \hat\sigma_{ie}e^{i\delta\varphi})e^{-\frac{i}{\hbar}\hat H_{atom}t} e^{i\Omega_b\hat b^+\hat b t}(\hat b - \hat b^+)e^{-i\Omega_b\hat b^+\hat b t}. \end{aligned} \tag{30}$$

At first consider the part containing the atom operators. As the matrix corresponding to $\hat H_{atom}$ is diagonal (see (5), (27)), then we obtain the following expression

$$exp(\frac{i}{\hbar}\hat H_{atom}t) = \begin{pmatrix} e^{i\omega_e t} & 0 & 0 \\ 0 & e^{i\omega_i t} & 0 \\ 0 & 0 & e^{i\omega_g t} \end{pmatrix}. \tag{31}$$

Utilizing (30) in (31) with the aid of (8) one can easily obtain that

$$\begin{aligned} exp(\frac{i}{\hbar}\hat H_{atom}t)\hat\sigma_{ig}exp(-\frac{i}{\hbar}\hat H_{atom}t)\hat\sigma_{ig} &= \\ = \hat\sigma_{ig}e^{i(\omega_i-\omega_g)t} \end{aligned}$$

In the same way, taking into account (9), (10) and (11)

$$exp(\frac{i}{\hbar}\hat H_{atom}t)\hat\sigma_{gi}exp(-\frac{i}{\hbar}\hat H_{atom}t)\hat\sigma_{ig} = \hat\sigma_{gi}e^{-i(\omega_i-\omega_g)t},$$

$$exp(\frac{i}{\hbar}\hat H_{atom}t)\hat\sigma_{ei}e^{-i\delta\varphi}exp(-\frac{i}{\hbar}\hat H_{atom}t) = \hat\sigma_{ei}e^{i(\omega_e-\omega_i)t}e^{-i\delta\varphi},$$

$$exp(\frac{i}{\hbar}\hat H_{atom}t)\hat\sigma_{ie}e^{i\delta\varphi}exp(-\frac{i}{\hbar}\hat H_{atom}t)\hat\sigma_{ig} = \hat\sigma_{ig}e^{-i(\omega_e-\omega_i)t}e^{i\delta\varphi}.$$

Thus, the atomic part of the interaction hamiltonian (30) is expressed in the following form

$$\begin{aligned} & e^{\frac{i}{\hbar}\hat H_{atom}t}(\hat\sigma_{ig} - \hat\sigma_{gi})e^{-\frac{i}{\hbar}\hat H_{atom}t} + e^{\frac{i}{\hbar}\hat H_{atom}t}(\hat\sigma_{ei}e^{-i\delta\varphi} - \hat\sigma_{ie}e^{i\delta\varphi})e^{-\frac{i}{\hbar}\hat H_{atom}t} = \\ & = \hat\sigma_{ig}e^{i(\omega_i-\omega_g)}t - \hat\sigma_{gi}e^{-i(\omega_i-\omega_g)t} + \hat\sigma_{ei}e^{i(\omega_e-\omega_i)t}e^{-i\delta\varphi} - \hat\sigma_{ei}e^{-i(\omega_e-\omega_i)t}e^{i\delta\varphi}. \end{aligned} \tag{32}$$

Now let us calculate the field operators. The time dependence of the operators in the interaction representation is determined by the Heisenberg equations of motion

$$\frac{d}{dt}\hat f = \frac{i}{\hbar}[\hat H_{field}, \hat f] \equiv \frac{i}{\hbar}(\hat H_{filed}\hat f - \hat f\hat H_{filed}), \tag{33}$$

where $\hat f$ stands for $\hat a$, $\hat a^+$, $\hat b$ and $\hat b$. Inserting $\hat H_{field}$ (15) in (33) we obtain

$$\frac{d}{dt}\hat f = \frac{i}{\hbar}\Omega(\hat f^+\hat f\hat f - \hat f\hat f^+\hat f). \tag{34}$$

Because of the commutation relation (14) the equation (34) is transformed into

$$\frac{d}{dt}\hat f = -i\Omega\hat f^+. \tag{35}$$

In the same manner

$$\frac{d}{dt}\hat f^+ = i\Omega\hat f^+. \tag{36}$$

So the time dependance of field operators is defined from (35) and (36) and are equal

$$\hat{a}(t) = \hat{a}(0)e^{-i\Omega_a t},\ \hat{a}(t)^{+} = \hat{a}^{+}(0)e^{i\Omega_a t},\ \hat{b}(t) = \hat{b}(0)e^{-i\Omega_b t},\ \hat{b}(t)^{+} = \hat{b}^{+}(0)e^{i\Omega_b t}, \tag{37}$$

Thus, by substituting (32) and (37) into the expression (30) we obtain the atom-photon interactions Hamiltonian in the interaction representation

$$\hat{H}^{(I)}_{\mathbf{rE}} = \hbar g_a(\hat{\mathbf{R}})[\hat{\sigma}_{ig}e^{i(\omega_i-\omega_g)t} - \hat{\sigma}_{gi}e^{-i(\omega_i-\omega_g)t}](\hat{a}e^{-i\Omega_a t} - \hat{a}^{+}e^{\Omega_a t})+$$
$$+\hbar g_a(\hat{\mathbf{R}})[\hat{\sigma}_{ei}e^{i(\omega_e-\omega_i)t}e^{-i\delta\varphi} - \hat{\sigma}_{ie}e^{-i(\omega_e-\omega_i)t}e^{i\delta\varphi}](\hat{b}e^{-i\Omega_b t} - \hat{b}^{+}e^{\Omega_b t}).$$

Introduce the following notation

$$\omega_i - \omega_j = \omega_{ij}. \tag{38}$$

Taking into account (38) the Hamiltonian is given by the following expression

$$\hat{H}^{(I)}_{\mathbf{rE}} = \hbar g_a(\hat{\mathbf{R}})[\hat{\sigma}_{ig}e^{i\omega_{ig}t} - \hat{\sigma}_{gi}e^{-i\omega_{ig}t}](\hat{a}e^{-i\Omega_a t} - \hat{a}^{+}e^{i\Omega_a t})+$$
$$+\hbar g_b(\hat{\mathbf{R}})[\hat{\sigma}_{ei}e^{i\omega_{ei}t}e^{i\delta\varphi} - \hat{\sigma}_{ie}e^{-i\omega_{ei}t}](\hat{b}e^{-i\Omega_b t} - \hat{b}^{+}e^{i\Omega_b t}),$$

or

(39)

$$\hat{H}^{(I)}_{\mathbf{rE}} = \hbar g_a(\hat{\mathbf{R}})(\hat{\sigma}_{gi}\hat{a}^{+}e^{i(\Omega_a-\omega_{ig})t}+\hat{\sigma}_{ig}\hat{a}e^{-i(\Omega_a-\omega_{ig})t}-\hat{\sigma}_{gi}\hat{a}e^{-i(\Omega_a+\omega_{ig})t}-e^{-i\delta\varphi}\hat{\sigma}_{ig}\hat{a}^{+}e^{i(\Omega_a+\omega_{ig})t}+$$
$$+\hbar g_b(\hat{\mathbf{R}})(\hat{\sigma}_{ie}\hat{b}^{+}e^{i(\Omega_b-\omega_{ei})t+i\delta\varphi}+\hat{\sigma}_{ei}\hat{b}e^{-i(\Omega_b-\omega_{ei})t-i\delta\varphi}-\hat{\sigma}_{ie}\hat{b}e^{-i(\Omega_b+\omega_{ei})t+i\delta\varphi}-\hat{\sigma}_{ei}\hat{b}^{+}e^{i(\Omega_b+\omega_{ei})t-i\delta\varphi}.$$

Further transformations require choosing particular transition scheme (Fig. 1).

**L-configuration.** For this configuration (Fig.1)

$$\omega_g < \omega_i < \omega_e \Rightarrow \omega_{ig} > 0, \omega_{ei} > 0. \tag{40}$$

After introducing notations

$$\Delta_a \equiv \Omega_a - \omega_{ig}$$

and

$$\Delta_b \equiv \Omega_b - \omega_{ei}.$$

from (39) and (40) we obtain the interaction Hamiltonian for L-configuration $\hat{H}^{(I,L)}_{\mathbf{rE}}$

$$\hat{H}^{(I,L)}_{\mathbf{rE}} = \hbar g_a(\hat{\mathbf{R}})(\hat{\sigma}_{gi}\hat{a}^{+}e^{i\Delta_a t} + \hat{\sigma}_{ig}\hat{a}e^{-i\Delta_a t} - \hat{\sigma}_{gi}\hat{a}e^{-i(\Omega_a+\omega_{ig})t} - \hat{\sigma}_{ig}\hat{a}^{+}e^{i(\Omega_a+\omega_{ig})t})+$$
$$+\hbar g_b(\hat{\mathbf{R}})(\hat{\sigma}_{ie}\hat{b}^{+}e^{i\Delta_b t+i\delta\varphi} + \hat{\sigma}_{ei}\hat{b}e^{-i\Delta_b t-i\delta\varphi} - \hat{\sigma}_{ie}\hat{b}e^{-i(\Omega_b+\omega_{ei})t+i\delta\varphi} - \hat{\sigma}_{ei}\hat{b}^{+}e^{i(\Omega_b+\omega_{ei})t-i\delta\varphi}).$$

It is easy to see that operators $\hat{\sigma}_{gi}\hat{a}$, $\hat{\sigma}_{ig}\hat{a}^{+}$, $\hat{\sigma}_{ie}\hat{b}$ and $\hat{\sigma}^{+}_{ei}\hat{b}^{+}$ leading to the violation of energy conservation law are multiplied by factors containing the sums of field oscillating and atom transitions frequencies. Contrary, operators $\hat{\sigma}_{gi}\hat{a}^{+}$, $\hat{\sigma}_{ig}\hat{a}$, $\hat{\sigma}_{ie}\hat{b}^{+}$ and $\hat{\sigma}_{ei}\hat{b}$ are multiplied by factors containing the differences (detuning) of aforementioned frequencies $\Delta \equiv \Omega - \omega$. So in the interaction Hamiltonian we have two groups of terms. The terms of the first group oscillate approximately with double optical frequencies and their contribution in the Schrödinger equation is vanishingly small. The terms of another group oscillate with detuning frequencies $\Delta$, and vary slowly. So their contribution in the Schrödinger equation is significant. These terms are $\hat{\sigma}_{gi}\hat{a}^{+}$, $\hat{\sigma}_{ig}\hat{a}$, $\hat{\sigma}_{ie}\hat{b}^{+}$ and $\hat{\sigma}_{ei}\hat{b}$. So utilize the averaging procedure - neglecting the fast oscillating terms such as $\hat{\sigma}_{gi}\hat{a}$, $\hat{\sigma}_{ig}\hat{a}^{+}$, $\hat{\sigma}_{ie}\hat{b}$ and $\hat{\sigma}_{ei}\hat{b}^{+}$ the interaction Hamiltonian reduces to the following form

$$\hat{H}^{(I,L)}_{\mathbf{rE}} \cong \hat{H}^{L}_{int} = \hbar g_a(\hat{\sigma}_{gi}\hat{a}^{+}e^{i\Delta_a t} + \hat{\sigma}_{ig}\hat{a}e^{-i\Delta_a t}) + \hbar g_b(\hat{\sigma}_{ei}\hat{b}^{+}e^{i\Delta_b t+i\delta\varphi} + \hat{\sigma}_{ei}\hat{b}e^{-i\Delta_b t-i\delta\varphi}).$$

It is also possible to strongly validate given approach by the mathematical averaging, contrary to heuristic averaging used above. Also the high order contributions can be calculated in this case.

**Λ-configuration.** For this configuration (Fig.1)

$$\omega_g < \omega_i < \omega_e \Rightarrow \omega_{ig} > 0, \omega_{ei} < 0.$$

After introducing notations

$$\Delta_a \equiv \Omega_a - \omega_{ig}$$

and

$$\Delta_b \equiv \Omega_b - \omega_{ie},$$

we obtain the interaction Hamiltonian for $\Lambda$-configuration $\hat{H}^{(I,\Lambda)}_{\mathbf{rE}}$ in the interaction representation

$$(41)\quad \hat{H}^{(I,\Lambda)}_{\mathbf{rE}} = \hbar g_a(\hat{\mathbf{R}})(\hat{\sigma}_{gi}\hat{a}e^{i\Delta_a t} + \hat{\sigma}_{ig}\hat{a}e^{-i\Delta_a t} - \hat{\sigma}_{gi}\hat{a}e^{-i(\Omega_a+\omega_{ig})t} - \hat{\sigma}_{ig}\hat{a}^{+}e^{i(\Omega_a+\omega_{ig})t})+$$
$$+\hbar g_b(\hat{\mathbf{R}})(\hat{\sigma}_{ie}\hat{b}^{+}e^{i(\Omega_b+\omega_{ie})t+i\delta\varphi} + \hat{\sigma}_{ei}\hat{b}e^{-i\Delta_b t+i\delta\varphi} - \hat{\sigma}_{ei}\hat{b}e^{i\Delta_b t+i\delta\varphi} - \hat{\sigma}_{ei}\hat{b}^{+}e^{i\Delta_b t-i\delta\varphi}).$$

Averaging (41) by fast oscillations leads to

$$\hat{H}^{(I,\Lambda)}_{\mathbf{rE}} \cong \hat{H}^{\Lambda}_{int} = \hbar g_a(\hat{\sigma}^{-}_{gi}\hat{a}^{+}e^{i\Delta_a t} + \hat{\sigma}_{gi}\hat{a}e^{-i\Delta_a t}) + \hbar g_b(\hat{\sigma}^{-}_{ie}\hat{b}^{+}e^{-i\Delta_b t+i\delta\varphi} + \hat{\sigma}_{ei}\hat{b}e^{i\Delta_b t-i\delta\varphi}).$$

***$V$*-configuration.** For this configuration (Fig.1)

$$\omega_g < \omega_i < \omega_e \Rightarrow \omega_{ig} < 0, \omega_{ei} > 0.$$

After introducing notations

$$\Delta_a \equiv \Omega_a - \omega_{gi}$$

and

$$\Delta_b \equiv \Omega_b - \omega_{ei},$$

we obtain the interaction Hamiltonian for $V$-configuration $\hat{H}^{(I,V)}_{\mathbf{rE}}$ in the interaction representation

$$(42)\quad \hat{H}^{(I,\Lambda)}_{\mathbf{rE}} = \hbar g_a(\hat{\mathbf{R}})(\hat{\sigma}_{gi}\hat{a}^{+}e^{i(\Omega_a+\omega_{gi})t} + \hat{\sigma}_{ig}\hat{a}e^{-i(\Omega_a+\omega_{gi})t} - \hat{\sigma}_{gi}\hat{a}e^{-i\Delta_a t} - \hat{\sigma}_{ig}\hat{a}^{+}e^{i\Delta_a t}+$$
$$+\hbar g_b(\hat{\mathbf{R}})(\hat{\sigma}_{ie}\hat{b}^{+}e^{i\Delta_b t+i\delta\varphi} + \hat{\sigma}_{ei}\hat{b}e^{-i\Delta_b t-i\delta\varphi} - \hat{\sigma}_{ie}\hat{b}e^{-i(\Omega_b+\omega_{ei})t+i\delta\varphi} - \hat{\sigma}_{ei}\hat{b}^{+}e^{-i(\Omega_b+\omega_{ei})t-i\delta\varphi}).$$

Neglecting in (42) fast oscillating terms we have

$$\hat{H}^{(I,V)}_{\mathbf{rE}} \cong \hat{H}^{V}_{int} = -\hbar g_a(\hat{\sigma}_{gi}\hat{a}e^{-i\Delta_a t} + \hat{\sigma}_{gi}\hat{a}^{+}e^{i\Delta_a t}) + \hbar g_b(\hat{\sigma}_{ie}\hat{b}^{+}e^{i\Delta_b t+i\delta\varphi} + \hat{\sigma}_{ei}\hat{b}e^{-i\Delta_b t-i\delta\varphi}).$$

Therefore, in this section we have shown that in the RWA three-level atom interacting with two modes of standing wave resonator, in the interaction representation is described by the following Hamiltonians

L-configuration

$$(43)\quad \hat{H}^{(I,L)}_{\mathbf{rE}} \cong \hat{H}^{L}_{int} = \hbar g_a(\hat{\sigma}_{gi}\hat{a}^{+}e^{-i\Delta_a t} + \hat{\sigma}_{ig}\hat{a}e^{-i\Delta_a t}) + \hbar g_b(\hat{\sigma}_{ie}\hat{b}^{+}e^{i\Delta_b t+i\delta\varphi} + \hat{\sigma}_{ei}\hat{b}e^{-i\Delta_b t-i\delta\varphi}).$$

$\Lambda$-configuration

$$(44)\quad \hat{H}^{(I,\Lambda)}_{\mathbf{rE}} \cong \hat{H}^{\Lambda}_{int} = \hbar g_a(\hat{\sigma}_{gi}\hat{a}^{+}e^{i\Delta_a t} + \hat{\sigma}_{ig}\hat{a}e^{-i\Delta_a t}) + \hbar g_b(\hat{\sigma}_{ie}\hat{b}e^{-i\Delta_b t+i\delta\varphi} + \hat{\sigma}_{ei}\hat{b}e^{i\Delta_b^{+}t-i\delta\varphi}).$$

V-configuration

$$(45)\quad \hat{H}^{(I,V)}_{\mathbf{rE}} \cong \hat{H}^{V}_{int} = -\hbar g_a(\hat{\sigma}_{gi}\hat{a}e^{-i\Delta_a t} + \hat{\sigma}_{ig}\hat{a}^{+}e^{i\Delta_a t}) + \hbar g_b(\hat{\sigma}_{ie}\hat{b}^{+}e^{i\Delta_b t+i\delta\varphi} + \hat{\sigma}_{ei}\hat{b}e^{-i\Delta_b t-i\delta\varphi}).$$

Farther we will investigate three-level model concentrated on the internal degrees of freedom and neglecting center of mass movement.

## 4. THE QUANTUM DYNAMICS OF A THREE-LEVEL ATOM

In this section we consider the internal quantum dynamics of a three-level atom in the fields of two modes of standing wave resonator. This model is more complicated and less studied than the well-known two-level system (see [1], [2] and references cited there). So, consider the Schrödinger equation

$$(46)\qquad i\hbar\frac{d|\Psi(t)>}{dt} = \hat{H}_{int}|\Psi(t)>$$

for the state vector $|\Psi>$ in the interaction representation, where the interaction Hamiltonian $\hat{H}_{int}$ due to (43)-(45) contains the following terms

$$(47)\quad \hat{H}_{int} = \hat{H}^{L}_{int} = \hbar g_a(\hat{\sigma}^{-}_{gi}\hat{a}^{+}e^{i\Delta_a t} + \hat{\sigma}_{gi}\hat{a}e^{i\Delta_a t}) + \hbar g_b(\hat{\sigma}^{-}_{ie}\hat{b}^{+}e^{i\Delta_b t+i\delta\varphi} + \hat{\sigma}_{ei}\hat{b}e^{-i\Delta_b t-i\delta\varphi}),$$

$$(48)\quad \hat{H}_{int} = \hat{H}^{\Lambda}_{int} = \hbar g_a(\hat{\sigma}^{-}_{gi}\hat{a}^{+}e^{i\Delta_a t} + \hat{\sigma}_{gi}\hat{a}e^{-i\Delta_a t}) + \hbar g_b(\hat{\sigma}^{-}_{ie}\hat{b}^{+}e^{-i\Delta_b t+i\delta\varphi} + \hat{\sigma}_{ei}\hat{b}e^{i\Delta_b t-i\delta\varphi}),$$

$$(49)\quad \hat{H}_{int} = \hat{H}^{V}_{int} = -\hbar g_a(\hat{\sigma}^{-}_{gi}\hat{a}^{+}e^{-i\Delta_a t} + \hat{\sigma}_{gi}\hat{a}e^{i\Delta_a t}) + \hbar g_b(\hat{\sigma}^{-}_{ie}\hat{b}^{+}e^{i\Delta_b t+i\delta\varphi} + \hat{\sigma}_{ei}\hat{b}e^{-i\Delta_b t-i\delta\varphi}).$$

We will study the internal dynamics of the three-level system when the interaction Hamiltonian becomes time independent. It is the most convenient way to obtain the time evolution operator.

4.1. **Exact resonance in three-level atom.** In the case of exact resonance $\Delta = 0$ the interaction Hamiltonians (47)-(49) transform into the following ones

$$\hat{H}_{int} = \hat{H}^L_{int} = \hbar g_a(\hat{\sigma}_{gi}\hat{a}^+ + \hat{\sigma}_{ig}\hat{a}) + \hbar g_b(\hat{\sigma}_{ie}\hat{b}^+ e^{i\delta\varphi} + \hat{\sigma}_{ei}\hat{b}e^{-i\delta\varphi}), \tag{50}$$

$$\hat{H}_{int} = \hat{H}^\Lambda_{int} = \hbar g_a(\hat{\sigma}_{gi}\hat{a}^+ + \hat{\sigma}_{gi}\hat{a}) - \hbar g_b(\hat{\sigma}_{ie}\hat{b}e^{i\delta\varphi} + \hat{\sigma}_{ei}\hat{b}^+ e^{-i\delta\varphi}), \tag{51}$$

$$\hat{H}_{int} = \hat{H}^V_{int} = -\hbar g_a(\hat{\sigma}_{gi}\hat{a}^+ + \hat{\sigma}_{ig}\hat{a}^+) + \hbar g_b(\hat{\sigma}_{ie}\hat{b}^+ e^{i\delta\varphi} + \hat{\sigma}_{ei}\hat{b}e^{-i\delta\varphi}). \tag{52}$$

Now they do not contain the explicit time dependence and the Schrödinger equation (46) for (50)-(52) can be integrated formally to give

$$|\Psi(t) >= exp(-\frac{i}{\hbar}\hat{H}_{int}t)|\Psi(0) >,$$

where the unitary time evolution operator is defined as

$$\hat{U}(t, t_0 = 0) \equiv exp(-\frac{i}{\hbar}\hat{H}_{int}t) \tag{53}$$

Substituting in (53) the Hamiltonians (50)-(52), we obtain the time evolution operators for $L$-, $\Lambda$- and $V$-configurations of a three-level atom

$$\hat{U}^L(t, t_0 = 0) = \exp[-ig_a t(\hat{\sigma}_{gi}\hat{a}^+ + \hat{\sigma}_{ig}\hat{a}) - i\hbar g_b t(\hat{\sigma}_{ie}\hat{b}^+ e^{i\delta\varphi} + \hat{\sigma}_{ei}\hat{b}e^{-i\delta\varphi})], \tag{54}$$

$$\hat{U}^\Lambda(t, t_0 = 0) = \exp[-ig_a t(\hat{\sigma}_{gi}\hat{a}^+ + \hat{\sigma}_{ig}\hat{a}) - ig_b t(\hat{\sigma}_{ie}\hat{b}^+ e^{i\delta\varphi} + \hat{\sigma}_{ei}\hat{b}e^{-i\delta\varphi})], \tag{55}$$

$$\hat{U}^V(t, t_0 = 0) = \exp[ig_a t(\hat{\sigma}_{gi}\hat{a}^+ + \hat{\sigma}_{gi}\hat{a}) - i\hbar g_b t(\hat{\sigma}_{ie}\hat{b}^+ e^{i\delta\varphi} + \hat{\sigma}_{ei}\hat{b}e^{-i\delta\varphi})]. \tag{56}$$

The initial state vector

$$|\Psi(0) \equiv |\Psi_{field} > \otimes|\Psi_{atom} >= \tag{57}$$

$$\sum_{n_a=0}^{\infty}\sum_{n_b=0}^{\infty} w_{n_a}w_{n_b}[c_e|e> + c_i|i> + c_g|g>]|n>= \sum_{n_a=0}^{\infty}\sum_{n_b=0}^{\infty} w_{n_a}w_{n_b}\begin{pmatrix} c_e \\ c_i \\ c_g \end{pmatrix}|n_a>|n_b>,$$

represents the direct product of the field state vector

$$|\Psi_{field} >\equiv \sum_{n_a=0}^{\infty}\sum_{n_b=0}^{\infty} w_{n_a}w_{n_b}|n_a>|n_b> \tag{58}$$

and the atom state vector

$$|\Phi_{atom} >\equiv [c_e|e> + c_i|i> + c_g|g>]. \tag{59}$$

It is a convenient form for utilizing the unitary transformations of initial atom-field state vector. So, the dynamics of a three-level atom, in the case of exact resonance, is defined by the (54)-(56) time evolution operators. In the next section we calculate these operators and utilize the unitary transformations important for applications of quantum computation.

## 5. CALCULATION OF THE TIME EVOLUTION OPERATOR AND UNITARY TRANSFORMATIONS

5.1. **Calculation of the time evolution operator in terms of the exponential matrix.** Based on relations (8)-(10) let us rewrite the time evolution operators (54)-(56) in the matrix form

$$\hat{U}^{(L)}(t,t_0=0)=e^{-i\begin{pmatrix} 0 & e^{-i\delta\varphi}g_b t\hat{b} & 0 \\ e^{i\delta\varphi}g_b t\hat{b}^+ & 0 & g_a t\hat{a} \\ 0 & g_a t\hat{a}^+ & 0 \end{pmatrix}} \tag{60}$$

$$\hat{U}^{(\Lambda)}(t,t_0=0)=e^{-i\begin{pmatrix} 0 & -e^{-i\delta\varphi}g_b t\hat{b}^+ & 0 \\ -e^{i\delta\varphi}g_b t\hat{b} & 0 & g_a t\hat{a} \\ 0 & g_a t\hat{a}^+ & 0 \end{pmatrix}}, \tag{61}$$

$$\hat{U}^{(V)}(t,t_0=0)=e^{-i\begin{pmatrix} 0 & e^{-i\delta\varphi}g_b t\hat{b} & 0 \\ e^{i\delta\varphi}g_b t\hat{b}^+ & 0 & -g_a t\hat{a}^+ \\ 0 & -g_a t\hat{a} & 0 \end{pmatrix}}, \tag{62}$$

or in the general form as

$$\hat{U}(t,t_0=0)=e^{-i\begin{pmatrix} 0 & \hat{B} & 0 \\ \hat{B}^+ & 0 & \hat{A} \\ 0 & \hat{A}^+ & 0 \end{pmatrix}},$$

where

| | configurations: | $L$ | $\Lambda$ | $V$ |
|---|---|---|---|---|
| $\hat{A}$ | $=$ | $g_a t\hat{a}$ | $g_a t\hat{a}$ | $-g_a t\hat{a}^+$ |
| $\hat{A}^+$ | $=$ | $g_a t\hat{a}^+$ | $g_a t\hat{a}^+$ | $-g_a t\hat{a}$ |
| $\hat{B}$ | $=$ | $e^{-i\delta\varphi}g_b t\hat{b}$ | $-e^{i\delta\varphi}g_b t\hat{b}^+$ | $e^{-i\delta\varphi}g_b t\hat{b}$ |
| $\hat{B}^+$ | $=$ | $e^{i\delta\varphi}g_b t\hat{b}^+$ | $-e^{-i\delta\varphi}g_b t\hat{b}^+$ | $e^{i\delta\varphi}g_b t\hat{b}^+$ |

corresponding operators from (60)-(62), including the corresponding scalar factors for convenience.

5.2. **Semiclassical approach.** At the given stage we substantially simplify the situation by means of simplifying the field operators. Right hand sides of equations (37) include only $\hat{f}\exp i\Omega t$ type terms, which describe contribution due to free evolution of the field. On the other hand it means that the applied fields are sufficiently intense compared with what can be contributed by the atomic transitions. The field operators at time $t$ may then be replaced by their averages in the initial state [3]. We assume the fields to be in the coherent states (the field of laser radiation) $|\{\alpha_n\}>$ and $|\{\beta_n\}>$ $(|\alpha|,|\beta|>>1)$, replace the operators $\hat{a}$ $(t_0=0)$ and $b$ $(t_0=0)$ by $\alpha$ and $\beta$ in (60)-(62) and receive the following

$L$-configuration

$$\hat{U}^{(L)}(t,t_0=0)=e^{-i\begin{pmatrix} 0 & e^{-i\delta\varphi}g_\beta t\beta & 0 \\ e^{i\delta\varphi}g_\beta t\beta^* & 0 & g_a t\alpha \\ 0 & g_a t\alpha^* & 0 \end{pmatrix}},$$

$\Lambda$-configuration

$$\hat{U}^{(\Lambda)}(t,t_0=0)=e^{-i\begin{pmatrix} 0 & -e^{-i\delta\varphi}g_\beta t\beta^* & 0 \\ -e^{i\delta\varphi}g_\beta t\beta & 0 & g_\alpha t\alpha \\ 0 & g_\alpha t\alpha^* & 0 \end{pmatrix}},$$

$V$-configuration

$$\hat{U}^{(V)}(t,t_0=0)=e^{-i\begin{pmatrix} 0 & e^{-i\delta\varphi}g_\beta t\beta & 0 \\ e^{i\delta\varphi}g_\beta t\beta^* & 0 & -g_\alpha t\alpha^* \\ 0 & -g_\alpha t\alpha & 0 \end{pmatrix}}.$$

In general we have

$$\hat{U}(t,t_0=0)=e^{-i\begin{pmatrix} 0 & B & 0 \\ B^* & 0 & A \\ 0 & A^* & 0 \end{pmatrix}}, \tag{63}$$

where

$$\begin{array}{cccccc} & & \texttt{configurations:} & L & \Lambda & V \\ & & & & & \\ & A & = & g_\alpha t\alpha & g_\alpha t\alpha & -g_\alpha t\alpha^* \\ (64) & A^* & = & g_\alpha ta^* & g_\alpha ta^* & -g_\alpha t\alpha \\ & B & = & e^{-i\delta\varphi}g_\beta t\beta & -e^{-i\delta\varphi}g_\beta t\beta^* & e^{-i\delta\varphi}g_\beta t\beta \\ & B^* & = & e^{i\delta\varphi}g_\beta t\beta^* & -e^{-i\delta\varphi}g_\beta t\beta^* & e^{i\delta\varphi}g_\beta t\beta^* \end{array}$$

The exponential matrix (63) can be calculated to give
(65)

$$\hat{U}(t,t_0=0)=\begin{pmatrix} \frac{AA^*+BB^*cos(\sqrt{BB^*+AA^*})}{BB^*+AA^*} & -\frac{iBsin(\sqrt{BB^*+AA^*})}{\sqrt{BB^*+AA^*}} & -\frac{BA(cos(\sqrt{BB^*+AA^*})-1)}{BB^*+AA^*} \\ -\frac{iB^*sin(\sqrt{BB^*+AA^*})}{\sqrt{BB^*+AA^*}} & cos(\sqrt{BB^*+AA^*}) & -\frac{iAsin(\sqrt{BB^*+AA^*})}{\sqrt{BB^*+AA^*}} \\ \frac{A^*B^*(cos(\sqrt{BB^*+AA^*})-1)}{BB^*+AA^*} & -\frac{iA^*sin(\sqrt{BB^*+AA^*})}{\sqrt{BB^*+AA^*}} & \frac{BB^*+AA^*cos(\sqrt{BB^*+AA^*})}{BB^*+AA^*} \end{pmatrix},$$

5.3. **Unitary transformations.** So, we found the time evolution operator for the three-level atom interacting with two modes of field in the coherent states. As the initial state vector is represented for the field in the arbitrary state $|\Psi_{field}>\equiv\sum_{n_a=0}^{\infty}\sum_{n_b=0}^{\infty}w_{n_a}w_{n_b}|n_a>|n_b>$, than it is essential to rewrite this state vector for two modes in coherent state. The probability that there are $n_a$ ($n_b$) photons at the initial coherent state $|\alpha>$ ($|\beta>$) is given by the Poisson distribution [4], i. e.

$$w_{n_a}=\frac{<n_a>^{n_a}\ e^{-<n_a>}}{n_a!},w_{n_b}=\frac{<n_b>^{n_b}\ e^{-<n_b>}}{n_b!}, \tag{66}$$

where $<n>=|\alpha|^2$ ($|\beta|^2$). Due to (66) we can now write the initial state of field (58) and atom-field system (57) in more suitable form

$$|\Psi_{field}>\equiv\sum_{n_a=0}^{\infty}\sum_{n_b=0}^{\infty}\frac{<n_a>^{n_a}\ e^{-<n_a>}}{n_a!}\frac{<n_b>^{n_b}\ e^{-<n_b>}}{n_b!}|n_a>|n_b>=|\alpha>|\beta>, \tag{67}$$

$$\begin{aligned} |\Psi(0)>=|\Psi_{field}>\otimes|\Psi_{atom}>= \qquad (68) \\ =\sum_{n_a=0}^{\infty}\sum_{n_b=0}^{\infty}\frac{<n_a>^{n_a}\ e^{-<n_a>}}{n_a!}\frac{<n_b>^{n_b}\ e^{-<n_b>}}{n_b!}[c_e|e>+c_i|i>+c_g|g>]|n_a>|n_b>= \\ =\sum_{n_a=0}^{\infty}\sum_{n_b=0}^{\infty}\frac{<n_a>^{n_a}\ e^{-<n_a>}}{n_a!}\frac{<n_b>^{n_b}\ e^{-<n_b>}}{n_b!}\begin{pmatrix} c_e \\ c_i \\ c_g \end{pmatrix}|a_a>|n_b>= \\ =[c_e|e>+c_i|i>+c_g|g>]|\alpha>|\beta>=\begin{pmatrix} c_e \\ c_i \\ c_g \end{pmatrix}|\alpha>|\beta>. \end{aligned}$$

Now applying (65) to (68) we obtain the total state vector $|\Psi(t)>$ of a three-level atom in the coherent field of two modes for arbitrary time $t>t_0=0$
(69)

$$|\Psi(t)>=\hat{U}(t,t_0=0)[c_e|e>+c_i|i>+c_g|g>]|\alpha>|\beta>=\hat{U}(t,t_0=0)\begin{pmatrix} c_e \\ c_i \\ c_g \end{pmatrix}|\alpha>|\beta>.$$

5.4. **Unitary transformations for L-configuration.** For L-configuration (59), (69) gives

$$|\Psi(t)>=\hat{U}^{(L)}(t,t_0=0)[c_e|e>+c_i|i>+c_g|g>]|\alpha>|\beta>=\hat{U}^{(L)}(t,t_0=0)\begin{pmatrix} c_e \\ c_i \\ c_g \end{pmatrix}|\alpha>|\beta>.$$

and due to (65) and (64) we obtain

$$\hat{U}^{(L)}(t,t_0=0)=\frac{1}{g_a^2|\alpha|^2+g_b^2|\beta|^2}\times\left\{\begin{pmatrix} g_a^2|\alpha|^2 & 0 & -e^{-i\delta\varphi}g_ag_b\beta\alpha \\ 0 & 0 & 0 \\ -e^{i\delta\varphi}g_ag_b\beta^*\alpha^* & 0 & g_b^2|\beta|^2 \end{pmatrix}+\right.$$

$$+cos(t\sqrt{g_a^2|\alpha|^2+g_b^2|\beta|^2})\begin{pmatrix} g_b^2|\beta|^2 & 0 & e^{-i\delta\varphi}g_ag_b\beta\alpha \\ 0 & g_a^2\alpha^2+g_b^2\beta^2 & 0 \\ e^{i\delta\varphi}g_ag_b\beta^*\alpha^* & 0 & g_a^2|\alpha|^2 \end{pmatrix}-$$

$$\left.-sin(t\sqrt{g_a^2|\alpha|^2+g_b^2|\beta|^2})\begin{pmatrix} 0 & e^{-i\delta\varphi}g_b\beta & 0 \\ e^{i\delta\varphi}g_b\beta^* & 0 & g_a\alpha \\ 0 & g_a\alpha^* & 0 \end{pmatrix}\right\}.$$

Thus, from the time evolution operator one can extract three formal parts: time independent, time dependent as $\cos(Gt)$ and time dependent as $\sin(Gt)$. For the unitary operator $\hat{U}^{(L)}(t,t_0=0)$ an initial moment $t=0$ we obtain

$$\hat{U^{(L)}}(t=0)=\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \tag{70}$$

and

$$\Psi(t=0)=\begin{pmatrix} c_e(0)|\alpha>|\beta> \\ c_i(0)|\alpha>|\beta> \\ c_g(0)|\alpha>|\beta> \end{pmatrix} \tag{71}$$

or

$$\Psi(t=0)=c_e(0)|e,\alpha,\beta>+c_i(0)|i,\alpha,\beta>+c_g|g,\alpha,\beta>, \tag{72}$$

where the following notations are introduced $|e>|\alpha>|\beta>\equiv|e,\alpha,\beta>$, $|i>|\alpha>|\beta>\equiv|i,\alpha,\beta>$ and $|g>|\alpha>|\beta>\equiv|g,\alpha,\beta>$ . (70)-(72) confirm the main feature of the interaction representation, that in the absence of interaction the state vectors do not change [5].

## 6. PROBABILITY AMPLITUDE METHOD

In this section, we present a different but equivalent method to solve for the evolution of the atom-field system described in interaction picture

$$i\hbar\frac{d|\Psi(t)>}{dt}=\hat{H}_{int}|\Psi(t)> \tag{73}$$

and based on the solutions of the probability amplitudes. The atom-field system is described by the interaction Hamiltonian (39)

$$\hat{H}^{(I)}_{\mathbf{rE}}=\hbar g_a(\hat{\sigma}_{gi}\hat{a}^+e^{i(\Omega_a-\omega_{ig})t}+\hat{\sigma}_{ig}\hat{a}e^{-i(\Omega_a-\omega_{ig})t}-\hat{\sigma}_{gi}\hat{a}e^{-i(\Omega_a+\omega_{ig})t}+\hat{\sigma}_{ig}\hat{a}^+e^{i(\Omega_a+\omega_{ig})t}+ \tag{74}$$

$$+\hbar g_b(\hat{\sigma}_{ie}\hat{b}^+e^{i(\Omega_b-\omega_{ei})t+i\delta\varphi}+\hat{\sigma}_{ei}\hat{b}e^{-i(\Omega_b-\omega_{ei})t-i\delta\varphi}-\hat{\sigma}_{ie}\hat{b}e^{-i(\Omega_b+\omega_{ei})t+i\delta\varphi}-\hat{\sigma}_{ei}\hat{b}^+e^{i(\Omega_b+\omega_{ei})t-i\delta\varphi}.$$

where the RWA should be utilized. Further calculations are based on amplitude method of QED [4] and are very similar for different configurations. So we perform the calculations only for L-configuration and then present the final result for all of them.

6.1. **L-Configuration.** In the RWA, the Hamiltonian (74), due to configuration (see Fig. 1), transforms into the ((47)-(49)) interaction operators. Especially for L-configuration

$$\hat{H}_{int} = \hat{H}^L_{int} = \hbar g_a(\hat{\sigma}_{gi}\hat{a}^+ e^{i\Delta_a t} + \hat{\sigma}_{ig}\hat{a}e^{-i\Delta_a t}) + \hbar g_b(\hat{\sigma}_{ie}\hat{b}^+ e^{i\Delta_b t+i\delta\varphi} + \hat{\sigma}_{ei}\hat{b}e^{-i\Delta_b t-i\delta\varphi}). \tag{75}$$

In the exact resonances case ($\Delta = 0$ ) (75) transforms into

$$\hat{H}^L_{int} = \hbar g_a(\hat{\sigma}_{gi}\hat{a}^+ + \hat{\sigma}_{ig}\hat{a}) + \hbar g_b(\hat{\sigma}_{ie}\hat{b}^+ e^{i\delta\varphi} + \hat{\sigma}_{ei}\hat{b}e^{-i\delta\varphi}). \tag{76}$$

Now acting on the wave vector $|e, n_a, n_b>$ by the operator (76) step by step we obtain

$$\hat{H}^L_{int}|e, n_a, n_b >= \hbar g_b e^{i\delta\varphi}\sqrt{n_b+1}|e, n_a, n_b >, \tag{77}$$

$$\hat{H}^L_{int}|e, n_a, n_b+1 >= \hbar g_a\sqrt{n_a+1}|g, n_a+1, n_b+1 > +\hbar g_b e^{-i\delta\varphi}\sqrt{n_b+1}|e, n_a, n_b >, \tag{78}$$

$$\hat{H}^L_{int}|g, n_a+1, n_b+1 >= \hbar g_a\sqrt{n_a+1}|i, n_a, n_b+1 >, \tag{79}$$

where $n_a$ and $n_b$ represent the number of $\hbar\Omega_a$ and $\hbar\Omega_b$ photons. From (79) it is also easy to see that for $n_a = 0$ and for arbitrary $n_b$ (let us $n_b + 1 = m$)

$$\hat{H}^L_{int}|g, 0, m >= 0. \tag{80}$$

Now with the aid of obtained amplitudes (77)-(80) one proceeds to solve the equation of motion (73) for state vector of total system $|\Psi(t)>$ , i. e.,

$$|\Psi(t) >= \sum_{n_a=0,n_b=0}^{\infty} \Psi_{e,n_a,n_b}(t)|e, n_a, n_b > +\Psi_{i,n_a,n_b+1}(t)|i, n_a, n_b+1 > + \tag{81}$$

$$+\Psi_{g,n_a+1,n_b+1}(t)|g, n_a+1, n_b+1 > +\Psi_{g,0,m}(t)|g, 0, m >$$

Now substitute (81) in (73) and then projecting the resulting equations onto $< e, n_a, n_b|$, $< i, n_a, n_b+1|$,$< g, n_a+1, n_b+1|$, $< g, 0, m|$ we can obtain the following system of equations:

$$\dot{\Psi}_{e,n_a,n_b}(t) = -ig_b e^{-i\delta\varphi}\sqrt{n_b+1}\Psi_{i,n_a,n_b+1}(t), \tag{82}$$

$$\dot{\Psi}_{i,n_a,n_b+1}(t) = -ig_b e^{i\delta\varphi}\sqrt{n_b+1}\Psi_{e,n_a,n_b}(t) - ig_a\sqrt{n_a+1}\Psi_{g,n_a+1,n_b+1}(t), \tag{83}$$

$$\dot{\Psi}_{g,n_a+1,n_b+1}(t) = -ig_a e^{i\delta\varphi}\sqrt{n_a+1}\Psi_{i,n_a,n_b+1}(t), \tag{84}$$

$$\dot{\Psi}_{g,0,m}(t) = 0. \tag{85}$$

To solve the system of equations (82)-(84) we should define the initial conditions first. Under initial conditions the state vector has the following form (see (67))

$$|\Psi(t=0) >= |\Psi_{atom} \otimes |\Psi_{field} >\equiv [c_e|e > +c_i|i > +c_g|g >] \otimes \sum_{n_a=0}^{\infty}\sum_{n_b=0}^{\infty} w_{n_a}w_{n_b}|n_a > |n_b >,$$

or in notations of (81)

$$|\Psi(t=0) >= \sum_{n_a,n_b=0}^{\infty} [w_{n_a,n_b}c_e|e, n_a, n_b > +w_{n_a,n_b+1}c_i|i, n_a, n_b+1 > + \tag{86}$$

$$+w_{n_a+1,n_b+1}c_g|g, n_a+1, n_b+1 >] + w_{0,m}c_g|g, 0, m >$$

where $w_{n_a,n_b} = w_{n_a}w_{n_b}$.

Comparing (81) for $t = 0$ and (86) we obtain the initial conditions

$$\Psi_{e,n_a,n_b}(t=0) = w_{n_a,n_b}c_e, \tag{87}$$

$$\Psi_{i,n_a,n_b+1}(t=0) = w_{n_a,n_b+1}c_i, \tag{88}$$

$$\Psi_{g,n_a+1,n_b+1}(t=0) = w_{n_a+1,n_b+1}c_g, \tag{89}$$

$$\Psi_{g,0,m}(t=0) = \Psi_{g,0,m}(t=0) = w_{0,m}c_g. \tag{90}$$

Now, let us solve the system of equations (82)-(84) under initial conditions defined by (87)-(90). Multiplying (82)-(84) by $e^{-st}$ and using the Laplace transformation of the probability amplitudes the system of equations transforms to the following:

$$e^{-st}\dot{\Psi}_{e,n_a,n_b}(t) = -ig_b\sqrt{n_b+1}e^{-st}e^{-i\delta\varphi}\Psi_{i,n_a,n_b+1}(t), \tag{91}$$

$$e^{-st}\dot{\Psi}_{i,n_a,n_b+1}(t) = -ig_b\sqrt{n_b+1}e^{-st}e^{i\delta\varphi} - ig_a e^{-st}\sqrt{n_a+1}\Psi_{g,n_a+1,n_b+1}(t), \tag{92}$$

$$e^{-st}\dot{\Psi}_{e,n_a,n_b}(t) = -ig_b\sqrt{n_a+1}e^{-st}\Psi_{i,n_a,n_b+1}(t). \tag{93}$$

Now integrating (91)-(93) by $t$ we obtain

$$s\overline{\Psi}_{e,n_a,n_b}(s) + ig_b e^{-i\delta\varphi}\sqrt{n_b+1\Psi_{e,n_a,n_b+1}(s)} = \Psi_{e,n_a,n_b}(t=0),$$

$$ig_b e^{i\delta\varphi}\sqrt{n_b+1\Psi_{e,n_a,n_b+1}(s)} + s\overline{\Psi}_{e,n_a,n_b+1}(s) + ig_b\sqrt{n_a+1\Psi_{g,n_a+1,n_b+1}(s)} = \Psi_{i,n_a,n_b+1}(t=0),$$

$$ig_a\sqrt{n_a+1\Psi_{i,n_a,n_b+1}(s)} + s\overline{\Psi}_{g,n_a+1,n_b+1}(s) = \Psi_{g,n_a+1,n_b+1}(t=0).$$

Here $\overline{\Psi}(s) = \int_0^\infty \Psi(t)e^{-st}dt$ is the Laplace transform of $\Psi(t)$. This system of equations we also can write in the following matrix form:

$$U\begin{pmatrix} \overline{\Psi}_{e,n_a,n_b}(s) \\ \overline{\Psi}_{i,n_a,n_b+1}(s) \\ \overline{\Psi}_{g,n_a+1,n_b+1}(s) \end{pmatrix} = \begin{pmatrix} w_{n_a,n_b}c_e \\ w_{n_a,n_b+1}c_i \\ w_{n_a+1,n_b+1}c_g \end{pmatrix},$$

where

$$U = \begin{pmatrix} s & ig_a e^{-i\delta\varphi}\sqrt{n_b+1} & 0 \\ ig_b e^{i\delta\varphi}\sqrt{n_b+1} & s & ig_a\sqrt{n_a+1} \\ 0 & ig_a\sqrt{n_a+1} & s \end{pmatrix}.$$

Now if we invoke the inverse Laplace transform we obtain the following expression for probability amplitudes

$$\begin{pmatrix} \Psi_{e,n_a,n_b}(t) \\ \Psi_{i,n_a,n_b+1}(t) \\ \Psi_{g,n_a+1,n_b+1}(t) \end{pmatrix} = \frac{1}{2\pi i}\int_C e^{st}U^{-1}(s)ds \begin{pmatrix} w_{n_a,n_b}c_e \\ w_{n_a,n_b+1}c_i \\ w_{n_a+1,n_b+1}c_g \end{pmatrix}, \tag{94}$$

where

$$U^{-1} = \frac{1}{s(s+i\lambda)(1-i\lambda)} \times \begin{pmatrix} s^2 + g_a^2(n_a+1) & -isg_a e^{-i\delta\varphi}\sqrt{n_b+1} & -g_b e^{-i\delta\varphi}\sqrt{n_b+1}g_a\sqrt{n_a+1} \\ -isg_a e^{-i\delta\varphi}\sqrt{n_b+1} & s^2 & -isg_a\sqrt{n_a+1} \\ -g_b e^{-i\delta\varphi}\sqrt{n_b+1}g_a\sqrt{n_a+1} & i - isg_a\sqrt{n_a+1} & s^2 + g_a^2(n_a+1) \end{pmatrix},$$

and $\lambda = \sqrt{g_a^2(n_a+1) + g_b^2(n_b+1)}$. The poles of (94) coincide with roots of the determinant $U$ and equal $0, i\lambda, -i\lambda$. From this, due to the Cauchy second integral theorem it follows

$$\Psi_{e,n_a,n_b}(t) = \left\{\left[cos(\lambda t) + \frac{g_a^2}{\lambda^2}(n_a+1)(1-cos(\lambda t))\right]\Psi_{i,n_a,n_b+1}(0) - \left[i\frac{g_b}{\lambda}e^{-i\delta\varphi}\sqrt{n_b+1}sin(\lambda t)\right]\Psi_{i,n_a,n_b+1}(0) - \right.$$
$$\left. - \left[\frac{g_b g_a}{\lambda^2}e^{-i\delta\varphi}\sqrt{n_b+1}\sqrt{n_a+1}(1-cos(\lambda t))\right]\Psi_{g,n_a+1,n_b+1}(0)\right\},$$

$$\Psi_{e,n_a,n_b+1}(t) = \left\{\left[\frac{g_a}{\lambda}e^{i\delta\varphi}\sqrt{n_b+1}sin(\lambda t)\right]\Psi_{e,n_a,n_b}(0) + cos(\lambda t)\right]\Psi_{i,n_a,n_b+1}(0) -$$
$$\left. - \left[i\frac{g_b}{\lambda}\sqrt{n_a+1}sin(\lambda t)\right]\Psi_{g,n_a+1,n_b+1}(0)\right\},$$

$$\Psi_{e,n_a,n_b}(t) = \left\{\left[\frac{g_b g_a}{\lambda_n^2}e^{-i\delta\varphi}\sqrt{n_b+1}\sqrt{n_a+1}(1-cos(\lambda t))\right]\Psi_{i,n_a,n_b}(0) - \left[i\frac{g_a}{\lambda}\sqrt{n_a+1}sin(\lambda t)\right]\Psi_{i,n_a,n_b+1}(0) - \right.$$
$$\left. - \left[cos(\lambda t) + \frac{g_a^2}{\lambda_n^2}(n_b+1)(1-cos(\lambda t))\right]\Psi_{g,n_a+1,n_b+1}(0)\right\}.$$

Amplitudes for the configurations $\Lambda$ and $V$ are obtained in similar manner.

The evolution operator

$$\left(\begin{array}{ccc} u_{11}(t) & u_{12}(t) & u_{13}(t) \\ u_{21}(t) & u_{22}(t) & u_{23}(t) \\ u_{31}(t) & u_{32}(t) & u_{33}(t) \end{array}\right)$$

for the $L, \Lambda, V$ configurations has the following form:

$L$-configuration, $\lambda = \sqrt{g_a^2(n_a+1) + g_b^2(n_b+1)}$.

$$u_{11}^L(t) = cos(\lambda t) + \frac{g_a^2}{\lambda^2}(n_a+1)(1 - cos(\lambda t)),$$

$$u_{12}^L(t) = -i\frac{g_b}{\lambda}e^{-i\delta\varphi}\sqrt{n_b+1}sin(\lambda t),$$

$$u_{13}^L(t) = \frac{g_b g_a}{\lambda^2}e^{-i\delta\varphi}\sqrt{n_b+1}\sqrt{n_a+1}(1 - cos(\lambda t)),$$

$$u_{21}^L(t) = -i\frac{g_b}{\lambda}e^{i\delta\varphi}\sqrt{n_b+1}sin(\lambda t),$$

$$u_{22}^L(t) = cos(\lambda t),$$

$$u_{23}^L(t) = -i\frac{g_b}{\lambda}e^{-i\delta\varphi}\sqrt{n_a+1}sin(\lambda t),$$

$$u_{31}^L(t) = \frac{g_b g_a}{\lambda^2}e^{-i\delta\varphi}\sqrt{n_b+1}\sqrt{n_a+1}(1 - cos(\lambda t)),$$

$$u_{32}^L(t) = -i\frac{g_b}{\lambda_n}e^{-i\delta\varphi}\sqrt{n_a+1}sin(\lambda t),$$

$$u_{33}^L(t) = cos(\lambda t) + \frac{g_a^2}{\lambda^2}(n_a+1)(1 - cos(\lambda t)).$$

$\Lambda$-configuration, $\lambda = \sqrt{g_a^2(n_a+1) + g_b^2 n_b^2}$.

$$u_{11}^\Lambda(t) = cos(\lambda t) + \frac{g_a^2}{\lambda^2}(n_a+1)(1 - cos(\lambda t)),$$

$$u_{12}^\Lambda(t) = i\frac{g_b}{\lambda}e^{-i\delta\varphi}\sqrt{n_b}sin(\lambda t),$$

$$u_{13}^\Lambda(t) = \frac{g_b g_a}{\lambda^2}e^{-i\delta\varphi}\sqrt{n_b}\sqrt{n_a+1}(1 - cos(\lambda t)),$$

$$u_{21}^\Lambda(t) = i\frac{g_b}{\lambda}e^{i\delta\varphi}\sqrt{n_b}sin(\lambda t),$$

$$u_{22}^\Lambda(t) = cos(\lambda t),$$

$$u_{23}^\Lambda(t) = -i\frac{g_b}{\lambda}\sqrt{n_a+1}sin(\lambda t),$$

$$u_{31}^\Lambda(t) = \frac{g_b g_a}{\lambda^2}e^{i\delta\varphi}\sqrt{n_b}\sqrt{n_a+1}(1 - cos(\lambda t)),$$

$$u_{32}^\Lambda(t) = -i\frac{g_a}{\lambda}\sqrt{n_a+1}sin(\lambda t),$$

$$u_{33}^\Lambda(t) = cos(\lambda t) + \frac{g_b^2}{\lambda^2}n_b(1 - cos(\lambda t)).$$

V-configuration, $\lambda = \sqrt{g_a^2 n_a + g_b^2(n_b+1)}$.

$$u_{11}^V(t) = cos(\lambda t) + \frac{g_a^2}{\lambda^2}(n_a)(1 - cos(\lambda t)),$$

$$u_{12}^V(t) = -i\frac{g_b}{\lambda}e^{-i\delta\varphi}\sqrt{n_b+1}sin(\lambda t),$$

$$u_{13}^V(t) = \frac{g_b g_a}{\lambda^2}e^{-i\delta\varphi}\sqrt{n_b+1}\sqrt{n_a}(1 - cos(\lambda t)),$$

$$u_{21}^V(t) = -i\frac{g_b}{\lambda}e^{i\delta\varphi}\sqrt{n_b+1}sin(\lambda t),$$

$$u_{22}^V(t) = cos(\lambda t),$$

$$u_{23}^V(t) = i\frac{g_a}{\lambda}\sqrt{n_a}sin(\lambda t),$$

$$u_{31}^V(t) = \frac{g_b g_a}{\lambda^2}e^{i\delta\varphi}\sqrt{n_b+1}\sqrt{n_a}(1 - cos(\lambda t)),$$

$$u_{32}^V(t) = i\frac{g_a}{\lambda}\sqrt{n_a}sin(\lambda t),$$

$$u_{33}^{V}(t) = cos(\lambda t) + \frac{g_b^2}{\lambda^2}(n_b + 1)(1 - cos(\lambda t)).$$

Under the assumptions described above we have constructed evolution operators in explicit form. Performed analysis has shown that unlike the case of methods of semiclassical analysis — where the atom is quantized whereas the field is classical, the completely quantum approach shows that in the absence of fields the atom still oscillates and this must be taken into account in quantum calculations. Moreover quantum processing may be performed on the level of single photons.

## References

[1] W.P.Schleich. Quantum optics in phase space. Wiley-VCH, Berlin, 2001.
[2] Allen L., Eberly J. H. Optical Resonance and Two–Level Atoms, Wiley, New York, 1975.
[3] Puri R.R. Mathematical methods of quantum optics. - Berlin: Springer, 2001, pp. XIII+285.
[4] Scully M.O. and Zubairy M.S. Quantum Optics. - Cambridge: University Press, 1997, pp. XXI+630.
[5] Cohen-Tannoudji C, Dupont-Roc J., Grynberg G. Atom-photon interactions. - Weinheim: Wiley-VCH, 2004, pp. XXII+656.
[6] L.Liu. Optical implementation of parallel fuzzu logic. Opt.Commun. 73(3), p.183-187, 1989.
[7] I.Zhang, L.Liu et al. Fuzzy reasoning morphological operators and their optical emplementation. App.Opt. v.36, N 11, p. 2328, 1997.
[8] M.L.Arestova, A.Iu. Bikovski. The method of realization of the optoelectronic scheme in bese many lavued logics. Quantum electronics. vol. 10, N 2, 1995.
[9] S.V.Iablonskiĭ. Introduction in discrete mathematics. Moscow, 384 p. 2003.
[10] E.Ercolessi, G.Marmo, G.Morandi, N.Mukunda. Geometry of mixed states and degeneracy structure of geometric phases for multi-valued quantum systems. ArXiv: quant-ph/0105007.
[11] J.I. Cirac, P.Zoller. Phys.Rev.Lett. 74, 4091, 1995.

Joint Institute for Nuclear Research, Dubna, Russia
*E-mail address*: `ggk@jinr.ru`

Institute of Cybernetics, Georgian Academy of Sciences, Tbilisi, Georgia
*E-mail address*: `z_melikishvili@posta.ge`